









- Computer Interfaces, a Review,” *Sensors*, vol. 12, no. 2, pp. 1211–1279, Jan. 2012.
- [8] R. Rupp, S. C. Kleih, R. Leeb, J. del R. Millan, A. Kübler, and G. R. Müller-Putz, “Brain–Computer Interfaces and Assistive Technology,” 2014, pp. 7–38.
- [9] R. Xu, N. Jiang, C. Lin, N. Mrachacz-Kersting, K. Dremstrup, and D. Farina, “Enhanced low-latency detection of motor intention from EEG for closed-loop brain-computer interface applications,” *IEEE Trans. Biomed. Eng.*, vol. 61, no. 2, pp. 288–96, Feb. 2014.
- [10] S. Aliakbaryhosseinabadi *et al.*, “Influence of attention alternation on movement-related cortical potentials in healthy individuals and stroke patients,” *Clin. Neurophysiol.*, vol. 128, no. 1, pp. 165–175, Jan. 2017.
- [11] R. Xu *et al.*, “Movement-related cortical potentials in paraplegic patients: abnormal patterns and considerations for BCI-rehabilitation,” *Front. Neuroeng.*, vol. 7, Aug. 2014.
- [12] Y. Gu, D. Farina, A. R. Murguialday, K. Dremstrup, and N. Birbaumer, “Comparison of movement related cortical potential in healthy people and amyotrophic lateral sclerosis patients,” *Front. Neurosci.*, vol. 7, 2013.
- [13] Ren Xu *et al.*, “A Closed-Loop Brain–Computer Interface Triggering an Active Ankle–Foot Orthosis for Inducing Cortical Neural Plasticity,” *IEEE Trans. Biomed. Eng.*, vol. 61, no. 7, pp. 2092–2101, Jul. 2014.
- [14] M. Jochumsen, I. K. Niazi, K. Dremstrup, and E. N. Kamavuako, “Detecting and classifying three different hand movement types through electroencephalography recordings for neurorehabilitation,” *Med. Biol. Eng. Comput.*, vol. 54, no. 10, pp. 1491–1501, Oct. 2016.
- [15] M. Jochumsen, I. Khan Niazi, D. Taylor, D. Farina, and K. Dremstrup, “Detecting and classifying movement-related cortical potentials associated with hand movements in healthy subjects and stroke patients from single-electrode, single-trial EEG,” *J. Neural Eng.*, vol. 12, no. 5, p. 056013, Oct. 2015.
- [16] A. Schwarz, P. Ofner, J. Pereira, A. I. Sburlea, and G. R. Müller-Putz, “Decoding natural reach-and-grasp actions from human EEG,” *J. Neural Eng.*, vol. 15, no. 1, p. 016005, Feb. 2018.
- [17] “The Amyotrophic Lateral Sclerosis Functional Rating Scale. Assessment of activities of daily living in patients with amyotrophic lateral sclerosis. The ALS CNTF treatment study (ACTS) phase I-II Study Group,” *Arch. Neurol.*, vol. 53, no. 2, pp. 141–7, Feb. 1996.
- [18] J. M. Cedarbaum *et al.*, “The ALSFRS-R: a revised ALS functional rating scale that incorporates assessments of respiratory function. BDNF ALS Study Group (Phase III),” *J. Neurol. Sci.*, vol. 169, no. 1–2, pp. 13–21, Oct. 1999.
- [19] N. Mrachacz-Kersting *et al.*, “The effect of type of afferent feedback timed with motor imagery on the induction of cortical plasticity,” *Brain Res.*, vol. 1674, pp. 91–100, Nov. 2017.
- [20] M. Jochumsen, I. K. Niazi, N. Mrachacz-Kersting, N. Jiang, D. Farina, and K. Dremstrup, “Comparison of spatial filters and features for the detection and classification of movement-related cortical potentials in healthy individuals and stroke patients,” *J. Neural Eng.*, vol. 12, no. 5, p. 056003, Oct. 2015.
- [21] Sheng-Hsiou Hsu, T. Mullen, Tzyy-Ping Jung, and G. Cauwenberghs, “Online recursive independent component analysis for real-time source separation of high-density EEG,” in *2014 36th Annual International Conference of the IEEE Engineering in Medicine and Biology Society*, 2014, pp. 3845–3848.
- [22] F. Karimi, J. Kofman, N. Mrachacz-Kersting, D. Farina, and N. Jiang, “Detection of Movement Related Cortical Potentials from EEG Using Constrained ICA for Brain-Computer Interface Applications,” *Front. Neurosci.*, vol. 11, Jun. 2017.